PCH BEYOND-USE DATING LIST[1]

Injections		
Insulin vials/prefilled pens/cartridges		Store in **fridge** before first use; after first use, store all types of insulin at **room temperature** for 28 days, **EXCEPT** • insulin detemir (Levemir®) 42 days • insulin degludec (Tresiba®) 56 days
Dalteparin 25,000 units/mL 3.8 mL multi-dose vial		14 days at **room temperature** after puncturing
Glucagon-like peptide-1 (GLP-1) receptor agonists prefilled pens		Store in **fridge** before first use; after first use store **at room temperature**: • semaglutide (Ozempic®) for 56 days • liraglutide (Victoza®) for 30 days
Lorazepam multi-dose vial		Store in **fridge** (2-8°C) until punctured. Waste remainder in the vial.
Testosterone cypionate 100 mg/mL 10 mL multi-dose vial		56 days at **room temperature** after puncturing
Tuberculin PPD (Tubersol®) for TST (Mantoux)		28 days **in fridge** after puncturing
Oral Preparations		
Antibiotic suspensions	amoxicillin, cloxacillin, cephalexin, cefprozil, cefixime, ciprofloxacin	14 days **in fridge** after reconstitution
	clindamycin, clarithromycin	14 days **at room temperature** after reconstitution
	azithromycin, amoxicillin & clavulanic acid, cefuroxime	10 days **in fridge** after reconstitution
Dabigatran capsules (in bottle)		120 days after opening
Nitroglycerin tablets		Manufacturer's expiry date **on original package**
Unit dose packaged oral medications		1 year or manufacturer's expiry date, whichever is earlier
Eye & Ear Preparations *(Use the manufacturer beyond-use date in the package insert or product monograph after opening. If not specified by the manufacturer or in the list below, use 60 days after opening)*		
Bimatoprost **0.03%**		60 days after opening for all brands **EXCEPT** 30 days for Sandoz
Brimonidine **0.15% and 0.2%**		60 days after opening for all brands **EXCEPT** 28 days for APO & JAMP
Brinzolamide/Brimonidine (Simbrinza®)		125 days after opening
Carboxymethylcellulose sodium 0.5% (Refresh Tears®)		90 days after opening
Cromoglycate (Cromolyn®)		28 days after opening
Dorzolamide		60 days after opening for all brands **EXCEPT** 40 days for MED & JAMP
Fluoroquinolones (e.g., gatifloxacin, moxifloxacin)		28 days after opening
Fucithalmic eye drops		28 days after opening
Latanoprost		Store in **fridge** until opened then 42 days after opening
Latanoprostene bunod (Vyzulta®)		56 days after opening
Loteprednol (Lotemax™)		28 days after opening
NSAID Products e.g. diclofenac, ketorolac, nepafenac		28 days after opening for all products and brands
Olopatadine 0.1% or 0.2%		60 days after opening for all brands **EXCEPT** 28 days for APO, ACT & JAMP brands
Retinol palmitate (Ocunox®) eye ointment		6 months after opening
Sodium hyaluronate (Hylo®) drops or gel		6 months after opening
Systane® **Ointment**		30 days after opening
Trehalose/Sodium Hyaluronate (Thealoz Duo®)		90 days after opening
Timolol		28 days after opening
Timolol combination products	Timolol + Brimonidine	60 days after opening for all brands **EXCEPT** 28 days for APO
	Timolol + Dorzolamide	60 days after opening for all brands **EXCEPT** 28 days for JAMP
	Timolol + Latanoprost	Store in **fridge** until opened, then 70 days after opening
	Timolol + Travoprost	120 days after opening
Tobramycin 0.3% (Tobrex®)		28 days after opening
Travoprost **0.003%** (Izba®)		125 days after opening
Topical Preparations *(manufacturer's expiry date unless listed below)*		
Calcipotriol Cream or Ointment (Dovonex®)		6 months after opening
Calcipotriol/ Betamethasone Ointment (Dovobet®)		1 year after opening
Calcipotriol/ Betamethasone Gel (Dovobet®)		6 months after opening
Fusidic acid cream (Taro-Fusidic acid)		28 days after opening
Fusidic acid ointment & combinations (Fucidin H®)		3 months after opening
Silver Sulfasalazine 1% Cream (Flamazine®)	20 & 50 g tube	**7 days** after opening
	500 g jar	**24 hours** after opening
Inhalers & Devices *(Use manufacturer's expiry date unless listed below. Use until device is empty or until the BUD, whichever is earliest)*		
Aclidinium (Tudorza Genuair®)		90 days after opening the pouch
Budesonide/glycopyrronium/formoterol (Breztri™ Aerosphere®)		120 actuations pack → 90 days after opening the pouch
Ellipta devices (e.g. Breo®, Anoro®, Incruse®, Arnuity®, Trelegy®)		42 days after opening the lid of the tray
Fluticasone/salmeterol (Advair Diskus®, Wixela® Inhub®)		30 days after opening the foil pouch
Formoterol/aclidinium (Duaklir Genuair®)		60 days after opening the pouch
Nebules (e.g. salbutamol, ipratropium)		Protect from light; 3 months once foil overwrap opened
Respimat devices (e.g. Spiriva®, Inspiolto®, Combivent®)		90 days after opening/puncture of the canister
Tiotropium (Spiriva®) 18 mcg capsule for inhalation		Expose 1 capsule at a time; if more than 1 capsule is exposed to air accidentally, the exposed unused capsule(s) must be discarded After first use, remaining capsules in the strip to be used within 5 days
Spacer device (Aerochamber®)		Replace every year or before if needed
Spiriva® Handihaler device		Replace every year or before if needed

[1]***Beyond-Use date**: the date after which a product must not be used assuming proper storage and hand hygiene practice.*
If the manufacturer specifies a beyond use date and it is different than listed above, the manufacturer's beyond use date is to be used